

$\pi^0 \rightarrow \gamma\gamma^*$, $\omega \rightarrow \pi^0\gamma^*$, and $\rho \rightarrow \pi\gamma^*$ decays in nuclear medium[★]

Agnieszka Bieniek, Anna Baran, Wojciech Broniowski

The H. Niewodniczański Institute of Nuclear Physics, PL-31342 Kraków, Poland

We calculate the medium modification of the $\pi\omega\rho$ vertex and analyze its significance for the $\pi^0 \rightarrow \gamma\gamma^*$, $\omega \rightarrow \pi^0\gamma^*$, and $\rho \rightarrow \pi\gamma^*$ decays in nuclear matter. We use a relativistic hadronic approach at zero temperature and in the leading-density limit, and consider decays of particles at rest with respect to the medium. It is shown that for the $\pi^0 \rightarrow \gamma\gamma^*$ the effects of the Δ isobar cancel almost exactly the effects of the nucleon ph excitations, such that the net medium effect is small. On the contrary, for the decays $\omega \rightarrow \pi^0\gamma^*$ and $\rho \rightarrow \pi\gamma^*$ we find a sizeable increase of the partial widths at virtualities of the photon in the range $0.3 - 0.6\text{GeV}$. The effect has direct significance for the calculation of dilepton yields from the Dalitz decays in relativistic heavy-ion collisions.

Keywords: Meson properties in nuclear medium, radiative and Dalitz meson decays in nuclear medium

PACS: 25.75.Dw, 21.65.+f, 14.40.-n

In recent years a lot of efforts have been undertaken in order to understand and quantitatively describe the properties of hadrons in nuclear matter. As an outcome, it has been commonly accepted that hadrons undergo substantial modifications induced by the medium [1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20], a view supported by numerous studies of two-point functions. Since the masses and widths of hadrons are significantly modified, one expects that also coupling constants are altered. However, there exist only a few investigations devoted to the issue of in-medium hadronic *three-point functions* [21,22,23,24,25,26,27]. In particular, it has been found that the $\rho\pi\pi$ coupling is largely increased in nuclear matter. This suggests the necessity of a careful examination of other couplings as well. In this paper we examine the *in-medium* $\pi\omega\rho$ vertex, which appears in such processes as $\pi^0 \rightarrow \gamma\gamma^*$, $\omega \rightarrow \pi^0\gamma^*$, and $\rho \rightarrow \pi\gamma^*$. These processes are important for theoretical studies of the dilepton production via

[★] Research supported by the Polish State Committee for Scientific Research grant 2P03B09419.

Dalitz decays in relativistic heavy-ion collisions. To our knowledge, the existing calculations and numerical codes use the vacuum value of the $\pi\omega\rho$ vertex. Clearly, medium effects, if present, alter the predictions of such calculations and this is the reason of our study. We find that for the Dalitz decays of vector mesons the presence of nuclear matter leads to a sizeable increase of partial widths: a factor of 2 for the ω and a factor of 3 for the ρ at the photon virtualities around 0.4GeV. This increase, in turn, directly leads to enhanced dilepton yields in processes $\omega \rightarrow \pi^0\gamma l\bar{l}$ and $\rho \rightarrow \pi\gamma l\bar{l}$, which may help to resolve the long-standing problem of the low-mass dilepton enhancement seen in relativistic heavy-ion collision [28,29].

In our analysis we apply a conventional hadronic model with meson, nucleon and Δ -isobar degrees of freedom, and work at the leading-density and zero temperature. We constrain ourselves to the situation where the decaying particle is at rest with respect to the medium. The extension to the general kinematic case is straightforward and will be presented elsewhere, together with the non-zero temperature analysis.

The vacuum value of the $\pi\omega\rho$ vertex (Feynman rule) is

$$-iV_{\pi\omega\rho}^{\nu\mu} = i\frac{g_{\pi\omega\rho}}{F_\pi}\epsilon^{\nu\mu pQ}, \quad (1)$$

where we have used the convenient short-hand notation $\epsilon^{\nu\mu pQ} = \epsilon^{\nu\mu\alpha\beta}p_\alpha Q_\beta$. We choose Q as the incoming momentum of the π , and p as the outgoing momentum of the ρ , and $q \equiv Q - p$ is the outgoing momentum of the ω . In a general case ω and ρ are a virtual particles. The value of $g_{\pi\omega\rho}$ is well known. For the case where all external momenta vanish, it can be obtained with help of the vector meson dominance from the anomalous $\pi\gamma\gamma$ coupling constant, namely $g_{\pi\omega\rho} = -\frac{g_\rho g_\omega}{e^2}g_{\pi\gamma\gamma}$, with $g_{\pi\gamma\gamma} = \frac{e^2}{4\pi^2 F_\pi}$ fixed by the anomaly.

Our calculation is made in the framework of a relativistic hadronic theory, where mesons interact with the nucleons and $\Delta(1232)$ isobars. The in-medium nucleon propagator can be conveniently decomposed into the *free* and *density* parts [4]:

$$S(k) = S_F(k) + S_D(k) = i(\gamma^\mu k_\mu + m_N)\left[\frac{1}{k^2 - m_N^2 + i\varepsilon} + \frac{i\pi}{E_k}\delta(k_0 - E_k)\theta(k_F - |k|)\right], \quad (2)$$

where m_N denotes the nucleon mass, $E_k = \sqrt{m_N^2 + k^2}$, and k_F is the Fermi momentum. The Rarita-Schwinger Δ propagator [30,31] has the form

$$iS_\Delta^{\mu\nu}(k) = i\frac{\gamma^\mu k_\mu + M_\Delta}{k^2 - M_\Delta^2}(-g^{\mu\nu} + \frac{1}{3}\gamma^\mu\gamma^\nu + \frac{2k^\mu k^\nu}{3M_\Delta^2} + \frac{\gamma^\mu k^\nu - \gamma^\nu k^\mu}{3M_\Delta}). \quad (3)$$




Fig. 1. Diagrams included in our calculation (crossed diagrams not shown). Wavy lines indicates the ρ , dashed lines the pions, solid lines the in-medium nucleon, and double lines the Δ .

We incorporate phenomenologically the effects of the finite width of the Δ by the replacement $M_\Delta \rightarrow M_\Delta - i\Gamma/2$.¹

The diagrams included in the calculation are displayed in Fig. 1, with the wavy lines denoting the ρ or ω , the dashed line the π^0 , the double line the Δ , and the single line the in-medium nucleon. Fermi-sea effects arise when one of the nucleon lines in each of the our diagrams of Fig. 1 involves the nucleon density propagator, S_D . For kinematic reasons, diagrams with more than one S_D vanish. Diagrams with no S_D propagator do not describe Fermi-sea effects; these vacuum-polarization diagrams are not considered in the present study.

The vertices (Feynman rules) needed for our calculation have the form

$$-iV_{\omega NN} = -ig_\omega \gamma^\mu, \quad (4)$$

$$-iV_{\rho NN} = i\frac{g_\rho}{2}(\gamma^\mu - \frac{i\kappa_\rho}{2m_N}\sigma^{\mu\nu}p_\nu)\tau^b, \quad (5)$$

$$-iV_{\pi NN} = \frac{g_A}{2F_\pi}Q\gamma^5\tau^a, \quad (6)$$

where p is the outgoing momentum of the virtual ρ , Q is incoming four-momentum of the π , and a , and b are the isospin indices of the π , and ρ respectively. For the interactions involving the Δ we have [31]

$$-iV_{N\Delta\pi} = g_{N\Delta\pi}Q^\alpha T^a, \quad (7)$$

$$-iV_{N\Delta\rho} = ig_{N\Delta\rho}(\not{p}\gamma^5 g^{\alpha\beta} - \gamma^\beta \gamma^5 p^\alpha)T^b, \quad (8)$$

$$-iV_{\Delta\Delta\omega} = -ig_\omega(\gamma^\alpha \gamma^\beta \gamma^\delta - \gamma^\delta g^{\alpha\beta} + \gamma^\alpha g^{\beta\delta} + \gamma^\beta g^{\delta\alpha}), \quad (9)$$

where T^a and T^b are the standard isospin $\frac{1}{2} \rightarrow \frac{3}{2}$ transition matrices [32]. Our choice of physical parameters is as follows:

¹ This treatment of the finite width of the Δ is consistent with the Ward-Takahashi identities for the $\pi\rho\omega$ vertex. This would not be true if Γ_Δ were introduced in the denominator of Eq. (3) only.

$$\begin{aligned}
g_\omega &= 10.4, & g_\rho &= 5.2, & g_A &= 1.26, & F_\pi &= 93\text{MeV}, \\
g_{N\Delta\pi} &= \frac{2.12}{m_\pi}, & g_{N\Delta\rho} &= \frac{2.12\sqrt{2}}{m_\pi}.
\end{aligned} \tag{10}$$

The values of the $N\Delta$ coupling constants follow from the non-relativistic reduction of the vertices and comparison to the non-relativistic values [32]. The $\Delta\Delta\omega$ coupling incorporates the principle of universal coupling. In the results presented below we have chosen $\kappa_\rho = 3.7$, *i.e.* $\kappa_\rho = \kappa_V$ value. Qualitatively similar conclusions follow when $\kappa_\rho = 6$ is used. One should note here that the relativistic form of couplings to the Δ , as well as the values of the coupling constants, are subject of an on-going discussion and research [31,33,34,35]. However, the ambiguities involved are not relevant for our study, where we explore the possibility and the size of the effect rather than seek accurate predictions.

In the following the results of the medium-modified vertex at the nuclear-saturation density ($\rho_0 = 0.17\text{fm}^{-3}$) will be presented relative to the vacuum value $g_{\text{vac}} = g_{\pi\rho\omega}$. We define

$$g_{\text{eff}} = g_{\pi\rho\omega} + \rho_0 B, \tag{11}$$

with B denoting the medium contribution evaluated according to the diagrams of Fig. 1. The in-medium $\pi\omega\rho$ vertex has the general tensor structure

$$\begin{aligned}
A^{\mu\nu} &= A_1 \varepsilon^{\mu\nu p Q} + A_2 \varepsilon^{\mu\nu u Q} + A_3 \varepsilon^{\mu\nu p u} + A_4 \varepsilon^{\mu\nu p Q} p^\nu \\
&+ A_5 \varepsilon^{\mu\nu p Q} Q + A_6 \varepsilon^{\mu\nu p Q} u^\nu + A_7 \varepsilon^{\nu\mu p Q} p^\mu + A_8 \varepsilon^{\nu\mu p Q} Q^\mu + A_9 \varepsilon^{\nu\mu p Q} u^\mu.
\end{aligned} \tag{12}$$

This structure, restricted by the Lorentz invariance and parity, is more general than in the vacuum case (1) due to the availability of the four-velocity of the medium, u . The result of *any* calculation will assume the form (12). The leading-density calculation of diagrams of Fig. 1 is very simple. As has been shown *e.g.* in Ref. [27], it amounts to evaluating the trace factors, substituting $k^\mu \rightarrow m_N u^\mu$, with $u^\mu = (1, 0, 0, 0)$ in the rest frame of the medium, and replacing the loop momentum integration by $\int \frac{d^4 k}{(2\pi)^4} \Theta(k_f - |\vec{k}|) \delta(k_0 - E_k) \rightarrow \frac{1}{8\pi} \rho_B$, where ρ_B is the baryon density. The vertex $A^{\mu\nu}$ satisfies the Ward-Takahashi identities: $q_\mu A^{\mu\nu} = 0$, and $p_\nu A^{\mu\nu} = 0$, which serves as a useful check of the algebra.

We begin the presentation of our numerical results with the process $\pi^0 \rightarrow \gamma\gamma$ (decay into two real photons), where the pion is *at rest* with respect to the medium. In this case the four-vectors Q and u are parallel, and out of the 9 structures in Eq. (12) only A_1 and A_3 survive: $A^{\mu\nu} = A_1 \varepsilon^{\mu\nu p Q} + A_3 \varepsilon^{\mu\nu p u}$. Thus in the rest frame of the medium, where $Q = (m_\pi, 0, 0, 0)$, we find $A^{\mu\nu} = A_1 Q_0 \varepsilon^{\mu\nu p 0} + A_3 u_0 \varepsilon^{\mu\nu p 0} = (m_\pi A_1 + A_3) \varepsilon^{\mu\nu p 0} = (A_1 + A_3/m_\pi) \varepsilon^{\mu\nu p Q}$. We combine the vacuum and medium pieces and introduce the notation $A^{\mu\nu} =$

$-i \frac{e^2}{g_\rho g_\omega} \left(\frac{g_{\pi\rho\omega}}{F_\pi} - \frac{\rho_B}{F_\pi} B \right) \epsilon^{\nu\mu p Q}$, which places the medium effects in the constant B . The contributions to B from diagrams (a), (b) and (c) of Fig.1 are as follows:

$$B_{(a)} = \frac{g_A g_\rho g_\omega (\kappa + 1)}{m_N (4m_N^2 - m_\pi^2)}, \quad (13)$$

$$B_{(b)} = \frac{g_{N\Delta\pi} g_{N\Delta\rho} g_\omega F_\pi m_\pi}{9M_\Delta^2 m_N [(m_N^2 - M_\Delta^2)^2 - m_\pi^2 m_N^2]} (m_N^4 + M_\Delta m_N^3 - 2M_\Delta^2 m_N^2 - (14)$$

$$- m_\pi^2 m_N^2 - M_\Delta^3 m_N + M_\Delta^4),$$

$$B_{(c)} = \frac{g_{N\Delta\pi} g_{N\Delta\rho} g_\omega F_\pi m_\pi}{27M_\Delta^4 [(m_N^2 - M_\Delta^2)^2 - m_\pi^2 m_N^2]} (-4m_N^5 - 6M_\Delta m_N^4 + 8M_\Delta^2 m_N^3 + (15)$$

$$+ 5M_\Delta^3 m_N^2 + (4m_N + 3M_\Delta) m_\pi^2 m_N^2 - 16M_\Delta^4 m_N - 11M_\Delta^5),$$

where M_Δ is understood to carry the width $i\Gamma_\Delta$. With parameters (10) we find numerically $B_{(a)} = 97\text{GeV}^{-3}$, $B_{(b)} = -3.1\text{GeV}^{-3}$, $B_{(c)} = -102\text{GeV}^{-3}$ for the formal case $\Gamma_\Delta = 0$. For the physical vacuum value of the Δ width, $\Gamma_\Delta = 0.12\text{GeV}$, we find $B_{(a)} = 97\text{GeV}^{-3}$, $B_{(b)} = -(2.6 + 1.5i)\text{GeV}^{-3}$, $B_{(c)} = -(87 + 42i)\text{GeV}^{-3}$ and $|g_{\text{eff}}/g_{\text{vac}}|^2 = |0.99 + 0.04i|^2 = 0.99$, whereas with nucleons only (diagram (a)) we would have $|g_{\text{eff}}^{(a)}/g_{\text{vac}}|^2 = 0.91^2 = 0.82$. Thus the effects of the Δ act in the opposite direction than the nucleons, resulting in an almost complete cancellation between diagrams (a) and (b,c) of Fig. 1.

The Dalitz decay of π^0 into the photon and the lepton pair proceeds through the decay into a real and a virtual photon, and subsequently the decay of the virtual photon into the lepton pair, $\pi^0 \rightarrow \gamma\gamma^* \rightarrow \gamma l\bar{l}$. The virtual photon can be either isovector (ρ -type) or isoscalar (ω -type). We denote the virtuality of γ^* as M , and investigate the dependence of g_{eff} on M . The results, displayed in Fig. 2, show that the change of M in the allowable kinematic range from 0 to m_π has a very little effect on the ratio of $|g_{\text{eff}}/g_{\text{vac}}|^2$. We notice a somewhat different behavior for the isoscalar and isovector photons at M approaching m_π , with the former ones coupled with greater strength. By comparing the curves corresponding to the calculation with the Δ (diagrams (a+b+c) of Fig. 1) and with nucleons only (diagrams (a)), we can see that the cancellation between diagrams (a) and (b+c) hold for the whole kinematically-available range. As a result, the medium effects on the Dalitz decays of π^0 are negligible, at most at the level of a few percent.

Next, we “turn around” the diagrams of Fig. 1 in order to consider the processes $\omega \rightarrow \pi^0\gamma^*$ and $\rho \rightarrow \pi^0\gamma^*$, where ω and ρ are on mass shell. In these cases we use the vacuum value $g_{\rho\omega\pi} = -1.13$, inferred from the experimental partial decay width of $\omega \rightarrow \pi^0\gamma$, rather than the value $g_{\pi\omega\rho} = -\frac{g_\rho g_\omega}{e^2} g_{\pi\gamma\gamma} = -1.36$ used earlier for the π^0 decay. The difference may be attributed to a form-factor effect: the virtuality of the vector meson is changed from 0 to the on-shell mass of the vector meson.




Fig. 2. The quantity $|g_{\text{eff}}/g_{\text{vac}}|^2$ for the process $\pi^0 \rightarrow \gamma^* \gamma$, plotted as a function of the virtual mass of γ^* , M , for the case $\rho_B = \rho_0$ and $\Gamma_\Delta = 120\text{MeV}$. The solid and dashed lines correspond to the case of isovector and isoscalar γ^* , respectively. The upper curves show the results of the full calculation, with diagrams (a,b,c) of Fig. 1 included, while the lower curves show the result of the calculation with diagram (a) only.

There is an interesting phenomenon in the decays of vector mesons, related to the analyticity of the amplitudes of Fig. 1 (b,c) in the virtuality of γ^* , denoted as M . For the case $\Gamma_\Delta = 0$ the amplitudes develop a pole at the location

$$M = \sqrt{\frac{m_v m_N^2 + m_\pi^2 m_N + m_v^2 m_N - M_\Delta^2 m_v + m_\pi^2 m_v}{m_N}} \approx 0.34\text{GeV}, \quad (16)$$

where $m_v = m_\rho$ or m_ω . The pole can be clearly seen in the plots of Fig. 3 for the case of $\Gamma_\Delta = 0$ (dashed lines). For the vacuum value of $\Gamma_\Delta = 120\text{MeV}$ (solid lines) the pole is washed-out, but its traces are still visible, with the curves reaching maxima around $M = 0.45\text{GeV}$. We observe that at low values of M the effective coupling constant remains unchanged for the ω decay, and is decreased for the ρ decay. However, at higher values of M , above 0.2GeV , the value of g_{eff} is larger than in the vacuum, with the effect quite large: an enhancement by a factor of 2 for the ω decay, and by a factor of 3 for the ρ decay. The exact numbers have to be treated with a grain of salt, since we have assumed the low-density limit, good as long as the results remain small. However, the values obtained are strongly indicative of possible large effects in the Dalitz decays of vector mesons in medium.

The enhancement of Fig. 3 will directly affect the calculations of the dilepton production in relativistic heavy-ion collisions, which is in fact very much desired. According to the results presented above, the theoretical yields from the Dalitz decays of vector mesons become considerably enhanced (a factor of 2 for the dominating ω meson) in the region of $0.4 - 0.5\text{GeV}$, which is precisely




Fig. 3. Top: $|g_{\text{eff}}/g_{\text{vac}}|^2$ for the decay $\omega \rightarrow \gamma^* \pi^0$ at the nuclear saturation density, plotted as a function of the virtuality of the photon, M . The dashed and solid lines correspond to the cases $\Gamma_\Delta = 0$ and 120 MeV, respectively. Bottom: the same for the decay $\rho \rightarrow \gamma^* \pi$.

where many existing calculations have problems in providing enough dileptons to explain the excess seen in the CERES and HELIOS experiments. At the same time, the Dalitz yields from the π^0 decays are not altered, which is also desired, as the theoretical calculations with the vacuum value of the $\pi\gamma\gamma^*$ vertex explain accurately the dilepton data in the low-mass (up to 0.2 GeV) region.

The authors are grateful to Wojciech Florkowski and Antoni Szczurek for numerous useful conversations.

References

- [1] G. E. Brown and M. Rho, Phys. Rev. Lett. **66** (1991) 2720
- [2] L. S. Celenza, A. Pantziris, C. M. Shakin, and W.-D. Sun, Phys. Rev. **C45** (1992) 2015
- [3] B. D. Serot and J. D. Walecka, Advances in Nuclear Physics **16** (1986) 1
- [4] S. A. Chin, Ann. Phys. (NY) **108** (1977) 301

- [5] T. Hatsuda, H. Shiomi, and H. Kuwabara, Prog. Theor. Phys. **95** (1996) 1009
- [6] H.-C. Jean, J. Piekarewicz, and A. G. Williams, Phys. Rev. **C49** (1994) 1981
- [7] M. Herrmann, B. L. Friman, and W. Noerenberg, Nucl. Phys. **A560** (1993) 411
- [8] B. Friman and H. J. Pirner, Nucl. Phys. **A617** (1997) 496
- [9] R. Rapp, G. Chanfray, and J. Wambach, Nucl. Phys. **A617** (1997) 472
- [10] S. Leupold and U. Mosel, Prog. Part. Nucl. Phys. **42** (1999) 221
- [11] S. Gao, C. Gale, C. Ernst, H. Stocker, and W. Greiner, Nucl. Phys. **A661** (1999) 518
- [12] T. Hatsuda and S. H. Lee, Phys. Rev. **C46** (1993) R34
- [13] S. H. Lee, Nucl. Phys. **A670** (2000) 119
- [14] U. Vogl and W. Weise, Prog. Part. Nucl. Phys. **27** (1991) 195
- [15] F. Klingl, N. Kaiser, and W. Weise, Nucl. Phys. **A624** (1997) 527
- [16] G. Chanfray, Nucl. Phys. **A685** (2001) 328
- [17] W. Broniowski and B. Hiller, Phys. Lett. **B392** (1997) 267
- [18] V. L. Eletsky, B. L. Ioffe, and J. I. Kapusta, Eur. J. Phys. **A3** (1998) 381
- [19] B. Friman, Acta Phys. Polon. **B29** (1998) 3195
- [20] M. Lutz, B. Friman, and G. Wolf, Nucl. Phys. **A661** (1999) 526
- [21] C. Song and V. Koch, Phys. Rev. **C54** (1996) 3218
- [22] W. Broniowski, W. Florkowski, and B. Hiller, Acta Phys. Polon. **B30** (1999) 1079
- [23] W. Broniowski, W. Florkowski, and B. Hiller, Eur. Phys. J. **A7** (2000) 287
- [24] M. Urban, M. Buballa, R. Rapp, and J. Wambach, Nucl. Phys. **A641** (1998) 433
- [25] M. Urban, M. Buballa, R. Rapp, and J. Wambach, Nucl. Phys. **A673** (2000) 357
- [26] M. K. Banerjee and J. A. Tjon, Phys. Rev. **C56** (1997) 497
- [27] W. Broniowski, W. Florkowski, and B. Hiller, nucl-th/0103027
- [28] CERES Collab., G. Agakichiev *et al.*, Phys. Rev. Lett. **75** (1995) 1272
- [29] HELIOS/3 Collab., M. Masera *et al.*, Nucl. Phys. **A590** (1995) 93c
- [30] W. Rarita and J. Schwinger, Phys. Rev. **60** (1941) 61
- [31] M. Benmerrouche, R. M. Davidson, and N. C. Mukhopadhyay, Phys. Rev. **C39** (1989) 2339

- [32] T. O. E. Ericson and W. Weise, *Pions and nuclei* (Clarendon Press, Oxford, 1988)
- [33] V. Pascalutsa and R. Timmermans, Phys. Rev. **C60** (1999) 042201
- [34] T. R. Hemmert, B. R. Holstein, and J. Kambor, J. Phys. G **G24** (1998) 1831
- [35] H. Haberzettl, preprint (1998), nucl-th/9812043